



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### HIV, violence, blame and shame

**Citation for published version:**

Pantelic, M, Boyes, M, Cluver, L & Meinck, F 2017, 'HIV, violence, blame and shame: Pathways of risk to internalized HIV stigma among South African adolescents living with HIV: Pathways', *Journal of the International AIDS Society*, vol. 20, no. 1, 21771. <https://doi.org/10.7448/IAS.20.1.21771>

**Digital Object Identifier (DOI):**

[10.7448/IAS.20.1.21771](https://doi.org/10.7448/IAS.20.1.21771)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Journal of the International AIDS Society

**Publisher Rights Statement:**

© 2017 Pantelic M et al; licensee International AIDS Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Research article

# HIV, violence, blame and shame: pathways of risk to internalized HIV stigma among South African adolescents living with HIV

Marija Pantelic<sup>§1,2</sup>, Mark Boyes<sup>1,3</sup>, Lucie Cluver<sup>1,4</sup> and Franziska Meinck<sup>1,5</sup>

<sup>§</sup>Corresponding author: Marija Pantelic, 32 Wellington Square, Oxford OX2 1ER, UK. Tel: +44 (0) 7442374711. ([marija.pantelic@spi.ox.ac.uk](mailto:marija.pantelic@spi.ox.ac.uk))

### Abstract

**Introduction:** Internalized HIV stigma is a key risk factor for negative outcomes amongst adolescents living with HIV (ALHIV), including non-adherence to anti-retroviral treatment, loss-to-follow-up and morbidity. This study tested a theoretical model of multi-level risk pathways to internalized HIV stigma among South African ALHIV.

**Methods:** From 2013 to 2015, a survey using total population sampling of ALHIV who had ever initiated anti-retroviral treatment (ART) in 53 public health facilities in the Eastern Cape, South Africa was conducted. Community-tracing ensured inclusion of ALHIV who were defaulting from ART or lost to follow-up. 90.1% of eligible ALHIV were interviewed ( $n = 1060$ , 55% female, mean age = 13.8, 21% living in rural locations). HIV stigma mechanisms (internalized, enacted, and anticipated), HIV-related disability, violence victimization (physical, emotional, sexual abuse, bullying victimization) were assessed using well-validated self-report measures. Structural equation modelling was used to test a theoretically informed model of risk pathways from HIV-related disability to internalized HIV stigma. The model controlled for age, gender and urban/rural address.

**Results:** Prevalence of internalized HIV stigma was 26.5%. As hypothesized, significant associations between internalized stigma and anticipated stigma, as well as depression were obtained. Unexpectedly, HIV-related disability, victimization, and enacted stigma were not directly associated with internalized stigma. Instead significant pathways were identified via anticipated HIV stigma and depression. The model fitted the data well (RMSEA = .023; CFI = .94; TLI = .95; WRMR = 1.070).

**Conclusions:** These findings highlight the complicated nature of internalized HIV stigma. Whilst it is seemingly a psychological process, indirect pathways suggest multi-level mechanisms leading to internalized HIV stigma. Findings suggest that protection from violence within homes, communities and schools may interrupt risk pathways from HIV-related health problems to psychological distress and internalized HIV stigma. This highlights the potential for interventions that do not explicitly target adolescents living with HIV but are sensitive to their needs.

**Keywords:** adolescent; stigma; abuse; shame; HIV/AIDS; structural equation modelling

To access the supplementary material to this article please see [Supplementary Files](#) under Article Tools online.

Received 20 December 2016; Accepted 8 August 2017; Published 21 August 2017

**Copyright:** © 2017 Pantelic M et al; licensee International AIDS Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Introduction

There are an estimated 2.1 million adolescents living with HIV (ALHIV) worldwide and 85% of them live in Sub-Saharan Africa [1]. South Africa is home to the world's largest population of ALHIV [2]. Globally, AIDS-related mortality is on the rise among ALHIV: between 2005 and 2015, AIDS-related mortality more than doubled in this high-risk population whilst all other age groups experienced reductions [3].

Internalized HIV stigma occurs when a person living with HIV internalizes perceived negative public attitudes towards people living with HIV and accepts them as applicable to

themselves [4]. By evoking strong feelings of shame and worthlessness [4], internalized HIV stigma can pose a serious threat to the long-term survival of ALHIV [5–7]. There are no quantitative studies on the prevalence or drivers of internalized HIV stigma among ALHIV in Sub-Saharan Africa [4]. In order to identify points for intervention for these adolescents, it is essential to understand mechanisms of risk for internalized HIV stigma in this population.

According to the HIV-stigma framework, internalized stigma is one of three core psychosocial HIV-stigma mechanisms experienced by people living with HIV [8,9]. Enacted HIV stigma refers to the extent to which people living with HIV are

discriminated against or treated differently based on their HIV status [10]. Anticipated HIV stigma refers to the extent to which people living with HIV anticipate negative public attitudes or differential treatment related to their HIV status [8]. The few Sub-Saharan African studies that tested associations between internalized HIV-stigma and other stigma mechanisms did this with adult samples, using simple bivariate correlations to assess validity of multidimensional HIV-stigma scales [4]. They found weak-to-moderate associations between enacted and internalized HIV stigma [10–12], and anticipated and internalized HIV stigma [13]. But these analyses were intended for psychometric scale assessments and hence did not take into account any potential confounding pathways or variables.

There is also a dearth of research on other potential drivers of internalized HIV stigma in Sub-Saharan Africa [4] – perhaps a reason for the lack of well-established programmes to address it [14]. A recent systematic review found that poor HIV-related health appeared to be a risk factor in three longitudinal studies, and depression appeared to be a risk factor in two longitudinal studies [4]. No other consistent predictors were found, and no quantitative studies assessing internalized HIV stigma among children or adolescents in the region were identified.

The literature on violence victimization and developmental psychology suggests that child maltreatment may also be a key driver of internalized HIV stigma via reduced psychological wellbeing [15,16]. Longitudinal findings from multiple systematic reviews consistently suggest that over time, child abuse victimization in the form of physical, sexual, emotional abuse and bullying leads to increases in depressive symptoms [17–22]. Similarly, enacted HIV stigma and bullying have been shown to have enduring negative impacts on depressive symptoms among youth from families affected by AIDS, which includes both ALHIV and HIV-negative adolescents [16].

Additionally, it is essential to assess whether and how HIV-related health is associated with maltreatment of ALHIV and internalized HIV stigma. Evidence from South Africa suggests that children from AIDS-affected families are at elevated risk of various forms of maltreatment when compared to their peers [16,23]. The maltreatment is not always exhibited as HIV-specific enacted stigma, but is nonetheless often indirectly tied to the children's health and HIV status. For example, ethnographic data from Brazil, South Africa, Zambia and Zimbabwe suggest that child abuse of AIDS-affected youth was linked to other household members' fear of HIV infection, and the perceived added burden of caring for ill children within a context of poverty [24–28]. AIDS-affected youths' experiences of abuse victimization were thus inextricably tied to their HIV-related health, and their perceptions of HIV stigma further shaped by abuse victimization. Whilst longitudinal research with adults has established a link between poor HIV-related health and internalized HIV stigma [4], qualitative research with ALHIV suggests that enacted HIV stigma and abuse victimization might account for this relationship [24–28]. However, these hypothesized pathways have not yet been tested quantitatively.

Taken together, this growing body of longitudinal research suggests likely pathways from HIV-related disability to internalized HIV stigma both directly [4], and via social and psychological risks [4,16,23]. Namely, HIV-related disability may place ALHIV at risk of more internalized HIV stigma via increased

likelihood of violence victimization and enacted HIV stigma (social risks), and via more anticipated HIV stigma and depression (psychological risks). However, neither of these potential pathways have been tested in structural equation models, nor explored among ALHIV.


This study aimed to develop and test a theoretical model of associations between HIV-related disability, hypothesized social (violence victimization and enacted HIV stigma) and psychological (anticipated HIV stigma and depression) risk factors and internalized HIV stigma. It draws on longitudinal quantitative findings from studies with adults living with HIV [4], qualitative research with AIDS affected adolescents in HIV-endemic contexts [24–28] and longitudinal research on psychological impacts of child maltreatment [17–21] to hypothesize pathways of risk (Figure 1).

## Methods

From 2013 to 2015, a survey of ALHIV was administered in the Eastern Cape, South Africa. First, all state healthcare facilities in two health districts were mapped. Facilities providing treatment for 5 or more ALHIV were included in the study ( $n = 53$ ). Second, all adolescents (ages 10–19) who had ever initiated ART in selected health facilities were sampled using clinic files. Patients were eligible if they met the 10–19 age bracket at the time of data collection, and if they were ever initiated on ART that was not short-term post-exposure prophylaxis. Second, adolescents ( $n = 1,176$ ) were traced in their communities so as to ensure inclusion of ALHIV who were not actively engaged in the healthcare system. One-on-one interviews ( $n = 1060$ ) were conducted in participants' homes, clinics or other places identified as safe and comfortable by the participants.

The questionnaire was pre-piloted with ALHIV, and a systematic review was conducted to ensure inclusion of key variables related to internalized HIV stigma into the questionnaire [4]. The questionnaire was developed in English, and items were translated and back translated independently by different Xhosa and English-speaking research assistants. Interviews lasted for about 90 min and were carried out on tablets. The full questionnaire consisted of nearly 450 items but tablet-assisted data collection allowed for skip patterns that reduced participant burden by omitting irrelevant questions. Participants chose whether they wanted to complete the interview directly on a mobile device (low-cost tablet), with the help of a research assistant, or a mix of the two depending on the questions. In case of low literacy levels or cognitive delay, Xhosa, English and Afrikaans-speaking interviewers, trained in working with ALHIV, read questions and provided assistance. Each completed questionnaire was assigned a serial number, which was recorded in the study roster. Data were linked to the serial number of their questionnaire and not participants' names resulting in an anonymized dataset. To ensure data protection from potential confidentiality breaches, data were accessible via password known only to the study management team.

Remuneration was not provided except for certificates, refreshments, and a toiletry pack. Voluntary informed consent was obtained from caregivers and adolescents. Where participants reported recent abuse, rape, suicidal attempt



**Figure 1. Hypothesized risk pathways from HIV-related disability to internalized HIV stigma.**

or other risk of significant harm, referrals were made to child protection and health services. Ethical clearance for this study was provided by the University of Oxford, University of Cape Town, South African National Departments of Health, Basic Education and Social Development as well as the Eastern Cape Departments of Health, Basic Education and Social Development.

## Measures

Age, gender and rural household location were recorded for descriptive purposes and for inclusion in the model as covariates.

*HIV-stigma mechanisms* were measured via the 10-item HIV-stigma scale for ALHIV (ALHIV-SS). The ALHIV-SS was developed in collaboration with ALHIV in South Africa and has been shown to have strong psychometric properties [29]. Enacted, anticipated and internalized HIV stigma were assessed via 3, 2 and 5 items, with Cronbach's  $\alpha$  levels of .57, .70 and .75 respectively. Latent variables of enacted, anticipated and internalized HIV stigma were used with individual items loaded onto them.

*HIV-related disability* was measured via two items asking about physical and cognitive disability adapted from The International Classification of Functioning, Disability and Health. The scale displayed good reliability (Cronbach's  $\alpha = .60$ ) [30]. Items were recoded into a dichotomous variable to capture any report of disability (0: no disability; 1: 1 or more disability).

*Violence victimization* was a latent measure with loaded factor scores for physical, emotional and sexual abuse, and bullying victimization. *Physical abuse* (2 items) and *emotional abuse* (10 items) were measured using items from the UNICEF Measures for National-level Monitoring

of Orphans and Other Vulnerable Children (Snider & Dawes, 2006). These measures have displayed strong reliability ( $\alpha = .70$ ) in previous studies of HIV/AIDS affected adolescents in South Africa [31]. Response options for all physical abuse and emotional abuse items were offered on a 5-point scale (0: never; 1: has happened, but not in the last year; 2: at least once this year; 3: monthly; 4: weekly). *Contact sexual abuse* was measured using three items from the Juvenile Victimization Questionnaire (JVQ) [32], also used in previous studies in South Africa [31]. *Bullying victimization* was measured with the 9-item 'Social and Health Assessment Peer Victimization Scale', used in previous studies with AIDS-affected children [16,33,34]. This scale was adapted from the Multidimensional Peer Victimization Scale, which was validated in the US [35]. Items include; being called names, being hit or threatened and having possessions broken or stolen. It demonstrated good internal consistency ( $\alpha = .79$ ) in the present sample.

*Depressive symptoms* were measured via the Child Depression Inventory short form (CDI-S), which has comparable results with the full CDI [36]. CDI-S has been used with AIDS-affected adolescents in South Africa, displaying acceptable internal consistency ( $\alpha = .67-.69$ ) [37]. CDI-S also demonstrated acceptable internal consistency in the present sample ( $\alpha = .62$ ). A latent variable was used in the present study, with individual items loading onto the depressive symptoms factor.

## Analysis strategy

The analysis was conducted in three stages using MPlus. First, to estimate the extent of bias in the sample, socio-demographic characteristics of eligible ALHIV who were not reached

were compared to the socio-demographic characteristics of ALHIV who were interviewed. These socio-demographic data were recorded from clinic files. Mean age between the included and excluded ALHIV was compared using z-scores. Gender and rurality frequencies between the included and excluded ALHIV were compared using  $\chi^2$ .

Second, prevalence rates of enacted, anticipated and internalized HIV stigma, HIV-related disability, and different types of violence victimization were calculated, and desegregated by gender. Dichotomous variables for different stigma mechanisms were defined to capture reports of any enacted, anticipated and internalized HIV stigma. Similarly, a dichotomous variable for any type of HIV-related disability was calculated. A dichotomous variable was created for *frequent emotional abuse* (0: once this year or less; 1: monthly or more often) and similarly for *frequent physical abuse* (0: once this year or less; 1: monthly or more often). Categorisations of frequent physical and emotional abuse were chosen based on previous studies with vulnerable children in South Africa, which aimed to clearly distinguish abuse from harsh parenting [38]. *Lifetime prevalence of contact sexual abuse* was also assessed (0: never sexually abused; 1: sexually abused). A dichotomous variable for bullying victimization above the mean was calculated. Mean scores for depressive symptoms were also calculated due to a lack of a validated clinical cutoff for the South African context.

Third, the structural equation model consisted of [1] a confirmatory factor analysis to confirm latent constructs and [2] a pathway model to assess a hypothesized theoretical model of internalized HIV-stigma risks (Figure 1). All included variables were latent constructs except for child disability, which was an observed variable. The model controlled for age, gender and rural household location. Model fit was assessed with: Comparative Fit Index (CFI) (>.90 indicates adequate fit), Tucker Lewis Index (TLI) (> .95 indicates adequate/good fit) [39–41], and Root Mean Square Error of Approximation (RMSEA) (< .05 indicated good model fit) [42].  $\chi^2$  was noted but not used to estimate model fit

because it is sensitive to sample size and is prone to Type 2 error [43,44].

## Results

90.1% ( $n = 1060$ ) of the eligible ALHIV were interviewed. 4.1% of ALHIV refused to participate (either caregiver or adolescent), 3.7% could not be traced, 0.9% were excluded due to severe cognitive delays and 1.2% were excluded due to other reasons such as emergency referrals, unsafe communities, and having moved out of study catchment area. The mean age of respondents was 13.8; 55.2% (587) of them were girls and 21.4% (228) lived in rural locations. A comparison of the included and excluded eligible samples based on known information: age, gender and rural/urban residential location identified no statistically significant differences between the interviewed and excluded samples (Table 1).

Table 2 provides further sample characteristics on the variables of interest, disaggregated by gender. Girls were more likely than boys to report any anticipated HIV stigma, frequent emotional abuse victimization and lifetime prevalence of sexual abuse victimization. No other gender differences were observed.

## Measurement model

The measurement model confirmed distinct latent constructs for enacted HIV stigma, violence victimization,

**Table 1. Comparisons between included and excluded ALHIV**

	HIV+ ( $n = 1060$ )	Excluded ( $n = 116$ )	Comparison tests
Age (mean, SD)	13.8, 2.834	14.8, 2.91	$p = .671$
Female ( $n$ , %)	587, 55.2%	66, 56.9%	$p = .769$
Rural ( $n$ , %)	228, 21.4%	26, 22.4%	$p = .813$

$p$ -Values associated with z score and  $\chi^2$  tests.

**Table 2. Sample characteristics**

	Boys living with HIV ( $n = 476$ )	Girls living with HIV ( $n = 584$ )	All ALHIV ( $n = 1060$ )	Gender comparison tests* ( $p$ value)
HIV-related disability	195, 41.0	238, 40.8	433, 40.8	.944
Internalized HIV stigma	103, 22.0	140, 25.3	243, 22.9	.206
Anticipated HIV stigma	116, 24.4	199, 34.1	315, 29.7	.001
Depression (mean, SD)	1.2, 1.8	1.3, 2.1	1.3, 2.0	.779
Enacted HIV stigma	31, 6.5	46, 7.9	77, 7.2	.400
Frequent physical abuse	28, 5.9	21, 3.6	49, 4.6	.078
Frequent emotional abuse	27, 5.7	52, 8.9	79, 7.5	.046
Lifetime prevalence of sexual abuse	17, 3.6	48, 8.2	65, 6.1	.002
Bullying victimization	162, 34.0	178, 30.5	340, 32.1	.218

\* $\chi^2$  and  $t$  tests were used to examine gender differences for dichotomous variables and scale variables, respectively.



anticipated HIV stigma, depression and internalized HIV stigma. Table 3 summarizes factor loadings of each indicator onto latent constructs of abuse victimization, enacted HIV stigma, anticipated HIV stigma, depression and internalized HIV stigma. The measurement model statistics indicated excellent model fit: RMSEA = .04; CFI = .984; TLI = .982; WRMR = .873;  $\chi^2(df) = 297.935 (242)$ .

### Structural model

The results of the final structural equation model are summarized in Figure 2. The model controlled for age, gender and rural household location. Rectangular shapes signify observed variables whereas ovals mark latent variables. Values indicate standardized  $\beta$  weights. Full lines indicate pathways that were significant at the  $p < .001$  level (\*\*\*). Dotted lines indicate hypothesized pathways that were

non-significant. HIV-related disability was hypothesized to have a direct effect on enacted HIV stigma, abuse victimization and internalized HIV stigma. HIV-related disability was hypothesized to predict enacted HIV stigma, abuse victimization and internalized HIV stigma. Enacted HIV stigma and abuse victimization were both hypothesized to be associated with depression, anticipated HIV stigma and internalized HIV stigma. Bidirectional associations between internalized HIV stigma and anticipated HIV stigma and depression were also hypothesized. Of these hypothesized associations, four were non-significant: (1) the direct association between HIV-related disability and internalized HIV stigma, (2) the direct association between abuse victimization and internalized HIV stigma, (3) the direct association between enacted HIV stigma and internalized HIV stigma and (4) the association between enacted HIV stigma and anticipated HIV stigma.

Internalized HIV stigma was directly associated with anticipated HIV stigma ( $\beta = .284, p < .001$ ), depressive symptoms ( $\beta = .445, p < .001$ ) and urban household location ( $\beta = -.014, p < .001$ ). Gender ( $\beta = .017, p = .196$ ) and age ( $\beta = -.011, p = .583$ ) were not significantly associated with internalized HIV stigma. In addition, the following pathways were identified. HIV-related disability was associated with more enacted HIV stigma ( $\beta = .493, p < .001$ ) and more abuse victimization ( $\beta = .372, p < .001$ ). Enacted HIV stigma was associated with more depressive symptoms ( $\beta = .294, p < .001$ ) but not with anticipated HIV stigma. Abuse victimization was associated with more anticipated HIV stigma ( $\beta = .334, p < .001$ ) and more depressive symptoms ( $\beta = .396, p < .001$ ). Indirect effects of HIV-related disability on higher depression scores via more abuse victimization ( $\beta = .147, p < .001$ ) and more enacted HIV stigma ( $\beta = .145, p < .001$ ) were observed. HIV-related disability was also indirectly associated with more anticipated HIV stigma via more abuse victimization ( $\beta = .124, p < .001$ ).

The fit of the final model was RMSEA = .023; CFI = .94; TLI = .95; WRMR = 1.070;  $\chi^2(df) = 3857.655 (372)$ . All fit statistics were excellent according to the pre-specified criteria.

### Discussion


This study aimed to test a theoretical model of hypothesized risk pathways to internalized HIV stigma among ALHIV in South Africa. As hypothesized, significant associations between internalized stigma and anticipated stigma, as well as depression were obtained. Unexpectedly, HIV-related disability, victimization, and enacted stigma were not directly associated with internalized stigma. These findings highlight the complicated nature of internalized HIV stigma. Whilst it is commonly seen as a purely psychological process, indirect pathways suggest multi-level mechanisms leading to internalized HIV stigma.

The present study expands on findings from longitudinal studies with adults living with HIV in the region, which found poor HIV-related physical health to be predictive of increases in internalized HIV stigma [4]. However, in the

**Table 3. Factor loadings for latent constructs**

	Standardized estimate
<b>Abuse victimization</b>	
Physical abuse	.458***
Emotional abuse	.659***
Sexual abuse	.407***
Bullying victimization	.648***
<b>Enacted HIV stigma</b>	
Stopped spending time with friends	.769***
Lost friends because of HIV	.674***
Teased because of HIV status	.990***
<b>Anticipated HIV stigma</b>	
People think that HIV-positive people are disgusting	.994***
People think that HIV is a punishment	.729***
<b>Depression</b>	
Personal outlook	.550***
Frequency of sadness	.695***
Feelings about appearance	.599***
Feelings towards self	.768***
Frequency of loneliness	.529***
Self-evaluation	.456***
Friends	.387***
Frequency of crying	.662***
Feelings of love	.623***
Bothered frequency	.769***
<b>Internalized HIV stigma</b>	
Does not feel as good as others because of their HIV status	.766***
Would rather die than live with HIV	.817***
Feels like a bad person for living with HIV	.833***
Feels ashamed of their HIV status	.780***
Feels dirty/contaminated inside because of HIV	.852***

\*\*\* indicates  $p < .001$



**Figure 2. Final structural equation model results.** Rectangular shape signifies an observed variable whereas ovals mark latent variables. Values indicate standardized  $\beta$  weights. Dotted lines indicate hypothesized pathways that were non-significant. Full lines indicate pathways that were significant. \*\*\* indicates  $p < .001$ ; \*\* indicates  $p < .005$ ; \* indicates  $p < .05$ . Model fit: RMSEA: .023; CFI: .94; TLI: .95; WRMR: 1.070. Model controlled for age, gender, rural household location.

present sample of ALHIV, HIV-related disability was associated with internalized HIV stigma only indirectly, via social risks (enacted HIV stigma and abuse victimization) and psychological risks (anticipated HIV stigma and depressive symptoms).

A growing corpus of research suggests that internalized HIV stigma hampers uptake of evidence-based HIV prevention, treatment and care [45–49] but most anti-stigma interventions target enacted HIV stigma [14]. There are only three published evaluations of interventions that target internalized HIV stigma in sub-Saharan Africa and these were delivered through clinics to adults living with HIV who were actively engaged in health care [50–52]. In light of present findings, a community-based approach to addressing internalized HIV stigma may be required for adolescents (Table 4). Our findings suggest that addressing individual-level risks – such as anticipated HIV stigma, depression and HIV-related health – will be essential to reduce internalized HIV stigma among ALHIV. However, present findings also suggest that tackling discrimination against ALHIV and violence victimization in homes, schools and communities may be essential to interrupt pathways of risk to internalized HIV stigma.

For example, interventions that do not necessarily target HIV-positive adolescents but are sensitive to their needs such as school-based anti-bullying programmes [53] might help reduce internalized HIV stigma. Parenting interventions specifically adapted for resource-limited settings might also help against internalized HIV stigma [54–56]. Youth violence exposure in one environment may impact involvement and/or victimization within another [57]. Therefore, violence

prevention programmes simultaneously targeting adolescents' local communities, homes and schools might be particularly beneficial.

Of note is that enacted HIV stigma was *not* associated with anticipated HIV stigma, which was one of the determinants of internalized HIV stigma. This is contrary to earlier studies, which focused on adults and used simple bivariate correlations [4]. Our findings suggest that HIV-related disability and/or violence victimization predict anticipated HIV stigma over and above enacted HIV stigma. Moreover, our findings suggest that enacted HIV stigma is associated with internalized HIV stigma only indirectly, via higher levels of depressive symptoms. More foundational and intervention research on the various HIV-stigma mechanisms is needed.

Gender differences were not observed for internalized or enacted HIV stigma in the initial descriptive analyses, and gender did not significantly contribute to the final model. Previous research in the region with adult samples generated inconsistent findings on the relationship between gender and internalized HIV stigma. For example, whilst two studies in Uganda and South Africa found that women were more likely to report internalized HIV stigma [58,59], another South African study found that women were less likely than men to report internalized HIV stigma [12]. To our knowledge, this is the first study in the region to look at this association within an adolescent, community-traced sample. Whilst findings challenge the common perception that girls are more likely to experience enacted HIV-related stigma [60–62], we also found that they were more likely to *anticipate* HIV stigma. More research on how gender is associated with different HIV-stigma mechanisms is clearly needed.

**Table 4. Recommendations for policy and practice**

**Policy makers**

1. Policies that do not specifically target ALHIV, but are sensitive to their needs might be beneficial. For example, evidence-based interventions that aim to reduce violence in schools, homes and communities may be beneficial for ALHIV even if they are not specifically designed for this population.
2. Evidence-based policies to support ALHIV and disabilities are urgently needed.

**Healthcare providers**

1. Taking into consideration the home and school environments of adolescent patients might help better support them.
2. Integrating psychological support and counselling into adolescent HIV care might help adolescents cope with HIV stigma.
3. In resource-limited contexts where psychosocial support is not available for all patients, it may be helpful to prioritize adolescents with physical or cognitive disabilities as they are at heightened risk of abuse and bullying victimization.

**Community organizations**

1. Family, school and community-based violence prevention programmes might help combat internalized HIV-related shame.
2. Healthcare providers may need support to better address the holistic needs of adolescent patients. Community organizations may help train healthcare providers to (a) routinely screen adolescent patients for mental health difficulties and history of violence victimization and (b) refer young people to adequate support services.
3. Teachers should also be supported and trained to detect bullying, discrimination and violence in schools, and refer adolescent students to adequate psychosocial support services.

**Schools**

1. Mechanisms to inform teachers of bullying and discrimination in schools are needed.
2. Violence prevention in schools that does not single out students living with HIV but rather targets more broadly peer and teacher violence may be beneficial for ALHIV.

This study has six important limitations that should be noted. First, the cross-sectional data limit inferences about directionality. More foundational, longitudinal research is needed to understand the mechanisms and their temporal sequencing. Second, all measures relied on self-report. Third, stigma is culturally embedded and socially constructed [63,64], and therefore HIV-stigma mechanisms might develop differently depending on the context. The present study reports on data from ALHIV in the Eastern Cape, South Africa and this may limit generalizability of findings. Fourth, this study did not examine protective factors. Evidence is urgently needed to better understand whether and how protective factors such as social support [4] may moderate pathways of risk to internalized HIV stigma. Fifth, only adolescents who had ever initiated ART were sampled. This may have limited reach of the most stigmatized adolescents who have not yet tested or initiated ART. However, unlike previous studies on internalized HIV stigma [4], this study community-traced adolescents in an attempt to reach most at-risk adolescents who had dropped out of treatment. Hence, adolescents who were missing clinic visits, defaulting or were lost to follow up were included in the study. Sixth, the length of interviews could have compromised data quality but inclusion of breaks and games sought to minimize participant burden.

This study also has a number of strengths. First, to our knowledge, this is the first study on internalized HIV stigma among ALHIV in Africa. Second, the structural equation model allowed the assessment of multiple inter-related risks. This enabled a more nuanced picture of how HIV-related disability – a key risk factor that had already been detected in previous research [4] – leads to internalized HIV stigma. Fourth, previous, adult-focused research on internalized HIV stigma in the region has largely overlooked interpersonal risk factors [4]. A key contribution of this study is that it empirically highlights that internalized HIV stigma is likely to be shaped by social contexts and power inequalities [65].

**Authors' affiliations**

<sup>1</sup>Department of Social Policy and Intervention, Oxford University, Oxford, UK; <sup>2</sup>International HIV/AIDS Alliance, Brighton, UK; <sup>3</sup>School of Psychology and Speech Pathology, Curtin University, Perth, Australia; <sup>4</sup>Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa; <sup>5</sup>OPTENTIA, School of Behavioural Sciences, North West University, Vanderbijlpark, South Africa

**Competing interests**

MP has no competing interests to declare. MB has no competing interests to declare. LC has no competing interests to declare. FM has no competing interests to declare.

**Authors' contribution**

MP conceptualized the paper, ran the analyses and drafted the manuscript. FM assisted with data analysis, interpretation of findings and manuscript write up. MB and LC assisted with study conceptualization, interpretation of findings and manuscript write up.

**Acknowledgements**

We would like to thank the participating adolescents who shared their time and stories with us, their families who welcomed us into their homes and the dedicated healthcare staff of facilities in the Eastern Cape. This study would not have been possible without the dedication, skill and benevolence of our research assistants who conducted these demanding interviews. MP is grateful to Dr. Elona Toska and Craig Carty for their ongoing support with this study.

**Funding**

This work was supported by the Nuffield Foundation under Grant [CPF/41513], the International AIDS Society through the Collaborative Initiative for Paediatric HIV Education and Research (CIPHER) grant (155-Hod), and the Oxford University Clarendon-Green Templeton College Scholarship (Marija Pantelic). Additional support for Lucie Cluver was provided by the European Research Council (ERC) under the European Union's Seventh Framework Programme (FP7/2007–2013)/ERC grant agreement n°313421. Additional funding was provided by Evidence for HIV Prevention in Southern Africa (EHPSA), a DFID programme managed by Mott MacDonald.

**References**

1. Idele P, Gillespie A, Porth T, Suzuki C, Mahy M, Kasedde S, et al. Epidemiology of HIV and AIDS among adolescents: current status, inequities, and data gaps. *J Acquir Immune Defic Syndr* [Internet]. United States: Idele, Priscilla. \*United Nations Children's Fund, New York, NY; and +Joint United Nations Programme on HIV/AIDS, Geneva, Switzerland;2014;66(Suppl 2): S144–53. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=medl&NEWS=N&AN=24918590>
2. World Health Organisation (WHO). Health for the world's adolescents. Geneva: WHO; 2014.
3. The United Nations Children's Fund (UNICEF). For every child, end AIDS: seventh stocktaking report. New York: UNICEF; 2016.
4. Pantelic M, Shenderovich Y, Cluver L, Boyes M. Predictors of internalised HIV-related stigma: a systematic review of studies in sub-Saharan Africa.



- Health Psychol Rev [Internet]. 2015 Jan;9(4):469–90. [cited 2016 Feb 8]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25559431>
5. Gross R, Yip B, Lo V, Iii R, Wood E, Alexander CS, et al. A simple, dynamic measure of antiretroviral therapy adherence predicts failure to maintain HIV-1 suppression. *J Infect Dis*. 2006;6021:1108–14.
6. Lima VD, Harrigan R, Bangsberg DR, Hogg RS, Gross R, Yip B, et al. The combined effect of modern highly active antiretroviral therapy regimens and adherence on mortality over time. *J Acquir Immune Defic Syndr*. 2009;50(5):529–36.
7. Reynolds SJ, Makumbi F, Nakigozi G, Kagaayid J, Graye RH, Wawere M, et al. HIV-1 transmission among HIV-1 discordant couples before and after the introduction of antiretroviral therapy. *AIDS*. 2012;25(4):473–77.
8. Earnshaw V, Chaudoir SR. From conceptualizing to measuring HIV stigma: a review of HIV stigma mechanism measures. *AIDS Behav* [Internet]. 2009 Dec;13(6):1160–77. [cited 2016 Feb 23]. Available from: [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4511707&tool=pmcentrez&render\\_type=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4511707&tool=pmcentrez&render_type=abstract)
9. Earnshaw V, Smith LR, Chaudoir SR, Amico KR, Copenhaver MM. HIV stigma mechanisms and well-being among PLWH: a test of the HIV stigma framework. *AIDS Behav* [Internet]. 2013 Jun;17(5):1785–95. [cited 2013 Dec 3]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23456594>
10. Holzemer WL, Uys LR, Chirwa ML, Greeff M, Makoe LN, Kohi TW, et al. Validation of the HIV/AIDS stigma instrument - PLWA (HASI-P). *AIDS Care* [Internet]. 2007 Sep;19(8):1002–12. [cited 2013 Oct 21]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17851997>
11. Kalichman SC, Simbayi LC, Cloete A, Mthembu PP, Mkhonta RN, Ginindza T. Measuring AIDS stigmas in people living with HIV/AIDS: the internalized AIDS-related stigma scale. *AIDS Care* [Internet]. 2009 Jan;21(1):87–93. [cited 2013 Oct 21]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19085224>
12. Simbayi LC, Kalichman S, Strebel A, Cloete A, Henda N, Mqeketo A. Internalized stigma, discrimination, and depression among men and women living with HIV/AIDS in Cape Town, South Africa. *Soc Sci Med* [Internet]. 2007 May;64(9):1823–31. [cited 2013 Oct 21]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17337318>
13. Visser M, Sipsma H. The experience of HIV-related stigma in South Africa. In: Liamputtong P, editor. *Stigma, discrimination and living with HIV/AIDS: a cross cultural perspective*. Dordrecht: Springer; 2013. p. 205–27.
14. Stangl AL, Lloyd JK, Brady LM, Holland CE, Baral S. A systematic review of interventions to reduce HIV-related stigma and discrimination from 2002 to 2013: how far have we come? *J Int AIDS Soc*. 2013;16:1–14.
15. Cluver L, Orkin M. Cumulative risk and AIDS-orphanhood: interactions of stigma, bullying and poverty on child mental health in South Africa. *Soc Sci Med* [Internet]. 2009 Oct;69(8):1186–93. [cited 2013 Feb]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19713022>
16. Boyes ME, Cluver LD. Relationships between familial HIV/AIDS and symptoms of anxiety and depression: the mediating effect of bullying victimization in a prospective sample of South African children and adolescents. *J Youth Adolesc* [Internet]. United States: Boyes, Mark E. Health Psychology and Behavioural Medicine Research Group, School of Psychology and Speech Pathology, Curtin University, GPO Box U1987, Perth, WA, 6845, Australia, mark.boyes@curtin.edu.au. 2015;44(4):847–59. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=prem&NEWS=N&AN=24996836>
17. Chen LP, Murad MH, Paras ML, Colbenson KM, Sattler AL, Goranson EN, et al. Sexual abuse and lifetime diagnosis of psychiatric disorders: systematic review and meta-analysis. *Mayo Clin Proc* [Internet]. 2010 Jul;85(7):618–29. [cited 2016 Mar 2]. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2894717&tool=pmcentrez&rendertype=abstract>
18. Hillberg T, Hamilton-Giachritsis C, Dixon L. Review of meta-analyses on the association between child sexual abuse and adult mental health difficulties: a systematic approach. *Trauma Violence Abuse* [Internet]. 2011 Jan;12(1):38–49. [cited 2016 May 30]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21288934>
19. Maniglio R. The impact of child sexual abuse on health: a systematic review of reviews. *Clin Psychol Rev* [Internet]. 2009 Nov;29(7):647–57. [cited 2016 Mar 28]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19733950>
20. Norman RE, Byambaa M, De R, Butchart A, Scott J, Vos T. The long-term health consequences of child physical abuse, emotional abuse, and neglect: a systematic review and meta-analysis. *PLoS Med* [Internet]. 2012 Jan;9(11):e1001349. [cited 2015 Apr 28]. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3507962&tool=pmcentrez&rendertype=abstract>
21. Pacheco JTB, Irigaray TQ, Memoriam BW (in, Nunes MLT, Argimon IIDL. Childhood maltreatment and psychological adjustment: a systematic review. *Psicol Reflexão e Crítica* [Internet]. 2014 Dec;27(4):815–24. [cited 2016 May 30]. Available from: [http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0102-79722014000400815&lng=en&nrm=iso&tlang=en](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0102-79722014000400815&lng=en&nrm=iso&tlang=en)
22. Ttofi MM, Farrington DP, Lösel F, Loeber R. Do the victims of school bullies tend to become depressed later in life? A systematic review and meta-analysis of longitudinal studies. Ttofi MM, editor. *J Aggress Confl Peace Res* [Internet]. 2011 May 16;3(2):63–73. [cited 2016 Dec 13]. doi:10.1108/17596591111132873.
23. Meinck F, Cluver LD, Boyes ME, Mhlomo EL. Risk and protective factors for physical and sexual abuse of children and adolescents in Africa: a review and implications for practice. *Trauma Violence Abuse* [Internet]. 2015 Jan;16(1):81–107. [cited 2016 Jun 13]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24648489>
24. Bond V. Stigma when there is no other option: understanding how poverty fuels discrimination toward people living with HIV in Zambia. In: Gillespie S, editor. *AIDS, poverty, and hunger: challenges and responses*. Washington (DC): International Food Policy Research Institute; 2006. p. 181–97.
25. Abadia-Barrero CE, Castro A. Experiences of stigma and access to HAART in children and adolescents living with HIV/AIDS in Brazil. *Soc Sci Med* [Internet]. 2006 Mar;62(5):1219–28. [cited 2014 Apr]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16099573>
26. Clay S, Bond V, Nyblade L. AIDS doesn't come through being together: children's experiences of HIV and AIDS related stigma in Zambia 2002-2003. Lusaka (Zambia): International HIV/AIDS Alliance and Pact; 2003.
27. Strode A, Grant KB. The role of stigma and discrimination in increasing the vulnerability of children and youth infected with and affected by HIV/AIDS. *Arcadia: Save the Children*; 2001.
28. Mahati ST, Chandiwana B, Munyati S, Chitiyo G, Mashange W, Chibatamoto P, et al. In: Munyati S, editor. *A qualitative assessment of orphans and vulnerable children in two Zimbabwean districts*. Cape Town (South Africa): HSRC Press; 2006.
29. Pantelic M, Boyes M, Cluver L, Thabang M. "They Say HIV is a Punishment from God or from Ancestors": cross-cultural adaptation and psychometric assessment of an HIV stigma scale for south african adolescents living with HIV (ALHIV-SS). *Child Indic Res* [Internet]. 2016 Nov 23. [cited 2016 Dec 8]. Available from: <http://link.springer.com/10.1007/s12187-016-9428-5>
30. WHO. International classification of functioning disability and health: ICF. Geneva (Switzerland): WHO; 2001.
31. Meinck F, Cluver LD, Boyes ME. Household illness, poverty and physical and emotional child abuse victimisation: findings from South Africa's first prospective cohort study. *BMC Public Health* [Internet]. 2015 Jan;15:444. [cited 2016 Jul 7]. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4418047&tool=pmcentrez&rendertype=abstract>
32. Finkelhor D, Hamby SL, Ormrod R, Turner H. The Juvenile Victimization Questionnaire: reliability, validity, and national norms. *Child Abuse Negl*. 2005;29:383–412.
33. Cluver L, Bowes L, Gardner F. Risk and protective factors for bullying victimization among AIDS-affected and vulnerable children in South Africa. *Child Abuse Negl* [Internet]. England: Cluver, Lucie. Department of Social Policy and Social Work, University of Oxford, UK. 2010;34(10):793–803. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med5&NEWS=N&AN=20880588>
34. Cluver L, Orkin M. Cumulative risk and AIDS-orphanhood: interactions of stigma, bullying and poverty on child mental health in South Africa. *Soc Sci Med* [Internet]. England: Cluver, Lucie. Department of Social Policy and Social Work, Oxford University, Oxford OX1 2ER, UK. lucie.cluver@socres.ox.ac.uk. 2009;69(8):1186–93. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med5&NEWS=N&AN=19713022>
35. Mynard H, Joseph S. Development of the multidimensional peer-victimization scale. *Aggress Behav*. 2000;26(2):169–78.
36. Kovacs M. *Manual: the children's depression inventory*. Toronto: Multi-Health Systems; 1995.
37. Cluver L, Orkin M, Boyes ME, Gardner F, Nikelo J. AIDS-orphanhood and caregiver HIV/AIDS sickness status: effects on psychological symptoms in South African youth. *J Pediatr Psychol* [Internet]. 2012 Sep;37(8):857–67. [cited 2012 Dec 29]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22313551>

38. Meinck F, Cluver LD, Boyes ME, Ndhlovu LD. Risk and protective factors for physical and emotional abuse victimisation amongst vulnerable children in South Africa. *Child Abuse Rev*. 2015;24(3):182–97.
39. Hu L, Bentler PM. Evaluating model fit. In: Hoyle R., editor: *Structural equation modeling: concepts, issues, and applications*. Thousand Oaks (CA): Sage; 1995. p. 76–99.
40. Bentler PM. Comparative fit indexes in structural models. *Psychol Bull*. 1990;107:238–46.
41. Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Struct Equ Model A Multidiscip J* [Internet]. 1999 Jan;6(1):1–55. doi:10.1080/10705519909540118
42. Bowen N, Guo S. *Structural equation modeling*. Oxford: Oxford University Press, Inc; 2012.
43. Schermelleh-Engel K, Moosbrugger H, Müller H. Evaluating the fit of structural equation models: tests of significance and descriptive goodness-of-fit measures. *Methods Psychol Res Online*. 2003;8(2):23–74.
44. Vandenberg RJ. Statistical and methodological myths and urban legends. *Organ Res Methods*. 2006;9(2):194–201.
45. Treves-Kagan S, El AM, Audrey A, Macphail C, Twine R, Maman S, et al. Gender, HIV testing and stigma: the association of HIV testing behaviors and community-level and individual-level stigma in rural South Africa differ for men and women. *AIDS Behav* [Internet]. 2017. doi:10.1007/s10461-016-1671-8.
46. Earnshaw VA, Smith LR, Shuper PA, Fisher WA, Cornman DH, Fisher JD, et al. HIV stigma and unprotected sex among PLWH in KwaZulu-Natal, South Africa: a longitudinal exploration of mediating mechanisms. *AIDS Care* [Internet]. 2014;26(12):1506–13. doi:10.1080/09540121.2014.938015.
47. Moshabela M, Zuma T, Orne-Gliemann J, Iwuji C, Larmarange J. “It is better to die”: experiences of traditional health practitioners within the HIV treatment as prevention trial communities in rural South Africa (ANRS 12249 TasP). *AIDS Care*. 2016;28(53):24–32.
48. Katz IT, Ryu AE, Onuegbu AG, Psaros C, Weiser SD, Bangsberg DR, et al. Impact of HIV-related stigma on treatment adherence: systematic review and meta-synthesis. *J Int AIDS Soc* [Internet]. 2013 Jan;16(3Suppl 2):18640. [cited 2013 Dec 3]. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3833107&tool=pmcentrez&rendertype=abstract>
49. Ng CK, Tsai AC. Proximate context of HIV-related stigma and women’s use of skilled childbirth services in Uganda. *AIDS Behav* [Internet]. 2017;21(1):307–16. doi:10.1007/s10461-016-1401-2.
50. Uys L, Chirwa M, Kohi T, Greeff M, Naidoo J, Makoe L, et al. Evaluation of a health setting-based stigma intervention in five African countries. *AIDS Patient Care STDS*. 2009;23(12):1059–66.
51. Tshabalala J, Visser M. Developing a cognitive behavioural therapy model to assist women to deal with HIV and stigma. *South African J Psychol* [Internet]. 2011 Mar 1;41(1):17–28. doi:10.1177/008124631104100103.
52. Maluccio JA, Wu F, Rokon RB, Rawat R, Wu F. Assessing the impact of food assistance on stigma among people living with HIV in Uganda using the HIV/AIDS stigma. *AIDS Behav* [Internet]. 2016;21(3):766–82. doi:10.1007/s10461-016-1476-9.
53. Cantone E, Piras AP, Vellante M, Preti A, Danielsdóttir S, D’Aloja E, et al. Interventions on bullying and cyberbullying in schools: a systematic review. *Clin Pract Epidemiol Ment Heal* [Internet]. 2015 Jan;11(Suppl1 M4):58–76. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4378064&tool=pmcentrez&rendertype=abstract>
54. Cluver L, Meinck F, Shenderovich Y, Ward CL, Romero RH, Redfern A, et al. A parenting programme to prevent abuse of adolescents in South Africa: study protocol for a randomised controlled trial. *Trials* [Internet]. 2016;1–10. doi:10.1186/s13063-016-1452-8.
55. Cluver L, Meinck F, Yakubovich A, Doubt J, Redfern A, Ward C, et al. Reducing child abuse amongst adolescents in low- and middle-income countries: a pre-post trial in South Africa. *BMC Public Health* [Internet]. 2016;16(567). doi:10.1186/s12889-016-3262-z.
56. Jordans MJD, Tol WA, Ndayisaba A, Komproe IH. A controlled evaluation of a brief parenting psychoeducation intervention in Burundi. *Soc Psychiatry Psychiatr Epidemiol*. 2013;48:1851–59.
57. Bowes L, Arseneault L, Arseneault L, Maughan B, Taylor A, et al. School, neighborhood, and family factors are associated with children’s bullying involvement: a nationally representative longitudinal study. *J Am Acad Child Adolesc Psychiatry*. 2009;48(5):545–53.
58. Sorsdahl K, Mall S, Stein DJ, Joska JA. The prevalence and predictors of stigma amongst people living with HIV/AIDS in the Western Province. *AIDS Care* [Internet]. 2011 Jun;23(6):680–85. [cited 2013 Oct 21]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21360358>
59. Nattabi B, Li J, Thompson SC, Orach CG, Earnest J. Factors associated with perceived stigma among people living with HIV/AIDS in post-conflict northern Uganda. *AIDS Educ Prev* [Internet]. 2011 Jun;23(3):193–205. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21696239>
60. Mahajan AP, Sayles JN, Patel VA, Remien RH, Szekeres G, Coates TJ. Stigma in the HIV/AIDS epidemic: a review of the literature and recommendations for the way forward. *AIDS*. 2010;22(Suppl 2):S67–79.
61. Campbell C, Gibbs A. Stigma, gender and HIV: case studies of intersectionality. In: Boesten J, Poku NK, editors. *Gender and HIV/AIDS: critical perspectives from the critical world*. Surrey (UK): Routledge; 2009. p. 29–46.
62. Osborne K, Tremlett L, Welbourn A, Hale F, Vasquez M. Piecing it together for women and girls: the gender dimensions of HIV-related stigma. London: International Planned Parenthood Federation; 2010.
63. Goffman E. *Stigma: the management of spoiled identity*. Harmondsworth: Penguin; 1968.
64. Kleinman A, Hall-Clifford R. Stigma: a social, cultural, and moral process. *J Epidemiol Community Health*. 2009;63(6):418–9.
65. Parker R, Aggleton P. HIV and AIDS-related stigma and discrimination: a conceptual framework and implications for action. *Soc Sci Med* [Internet]. 2003 Jul;57(1):13–24. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12753813>